
The future of dark matter search with

XENONINT









Adam Brown for the XENON collaboration

Liquid xenon TPCs

- ► Interaction ⇒ light (S1) & ionisation
- ► e- drift up to gas; e--Xe collisions in gas \implies light (S2)
- ► S2 hit pattern + drift time ⇒ 3D position
- S2/S1 ratio ⇒ type of interaction: electronic recoil (ER) / nuclear recoil (NR)






- XENONnT is an evolution of XENON1T, which set very strong limits on DM-nucleus interactions [1] and observed 0vDEC [2]
- Located in LNGS underground lab shielded against cosmic rays: rock shielding equivalent to 3600 m of water
- TPC contains 5.9 t xenon, three times more than XENON1T
- ► S1 and S2 readout with two arrays of 3" PMTs: in total 494
- All tested in liquid and gaseous xenon to check for stable operation and leaktightness





Radon reduction compared to XENON1T





Qdrive pumps

10.9 mBq

Neutron veto

- Neutrons are dangerous NR background
- New n-veto subsystem for XENONnT
- Inner region of existing muon veto
- Optically separate region, with 120 additional PMTs
- ► 0.5% Gd₂(SO₄)₃ in the water tank
- >80% efficiency tagging neutrons which have scattered once in the TPC





Liquid xenon purification

- Purification removes electronegative contamination so electrons reach the gas
- With gas purification ~1.8 t/day possible
- Liquid purification means 21 t/day


Porcupine


Crogenic system

1.9 mBq

2.4 mBq


Two custom cryogenic oxygen filters






Summary

- XENONnT is coming with unprecedented sensitivity to WIMP dark matter
- New technologies keep the large amount of xenon ultra-pure





- Construction is ongoing
- First data will be taken in 2020

References

- 1. Aprile et al., PRL 121 111302 (2018)
- 2. Aprile et al., Nature 568, 532 (2019)

cor

contact@xenon1t.org



www.xenon1t.org

